


Patterns

Anthroponumbers.org: A quantitative database of human impacts on Planet Earth

Graphical abstract



Authors

Griffin Chure, Rachel A. Banks, Avi I. Flamholz, ..., Yinon Bar-On, Ron Milo, Rob Phillips

Correspondence

griffinchure@gmail.com (G.C.), phillips@pboc.caltech.edu (R.P.)

In brief

The environmental impacts of human action on Earth are being felt on many fronts. Despite our deep knowledge of these impacts, finding reliable quantitative information is burdensome, often requiring domain expertise and programmatic acumen. We present the Human Impacts Database, which houses a diverse array of quantities regarding human impacts, making them easily accessible and searchable. We use this database to present a broad view of the Anthropocene, exploring the global magnitudes, spatial dependence, and temporal dynamics of human impacts.

Highlights

- We present a holistic view of the many ways humans alter Earth at a global scale
- We consider how these global quantities vary across geography
- We further explore the time- and population-dependent dynamics of these impacts
- We enumerate and describe key properties associated with each entry in the database



Descriptor

Anthroponumbers.org: A quantitative database of human impacts on Planet Earth

Griffin Chure,^{1,2,10,11,*} Rachel A. Banks,^{3,4,5,10} Avi I. Flamholz,^{3,4} Nicholas S. Sarai,⁶ Mason Kamb,⁵ Ignacio Lopez-Gomez,^{4,7} Yinon Bar-On,⁸ Ron Milo,⁸ and Rob Phillips^{3,5,9,*}

¹Department of Biology, Stanford University, Stanford, CA, USA

²Department of Applied Physics, California Institute of Technology, Pasadena, CA, USA

³Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, CA, USA

⁴Resnick Sustainability Institute, California Institute of Technology, Pasadena, CA, USA

⁵Chan-Zuckerberg BioHub, San Francisco, CA, USA

⁶Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, CA, USA

⁷Department of Environmental Science and Engineering, California Institute of Technology, Pasadena, CA, USA

⁸Department of Plant and Environmental Sciences, Weizmann Institute of Science, Rehovot, Israel

⁹Department of Physics, California Institute of Technology, Pasadena, CA, USA

¹⁰These authors contributed equally

¹¹Lead contact

*Correspondence: griffinchure@gmail.com (G.C.), phillips@pboc.caltech.edu (R.P.)

<https://doi.org/10.1016/j.patter.2022.100552>

THE BIGGER PICTURE Over the last 10,000 years, human activities have transformed Earth through farming, forestry, mining, and industry. The complex results of these activities are now observed and quantified as “human impacts” on Earth’s atmosphere, oceans, biosphere, and geochemistry. While myriad studies have explored facets of human impacts on the planet, they are necessarily technical and often highly focused. Thus, finding reliable quantitative information requires a significant investment of time to assess each quantity and associated uncertainty. We present the Human Impacts Database (www.anthroponumbers.org), which houses a diverse array of such quantities. We review a subset of these values and how they help build intuition for understanding the Earth-human system. While collation alone does not tell us how to best ameliorate human impacts, we contend that any future plans should be made in light of a quantitative understanding of the interconnected ways in which humans influence the planet.



Production: Data science output is well understood and (nearly) universally adopted

SUMMARY

The Human Impacts Database (www.anthroponumbers.org) is a curated, searchable resource housing quantitative data relating to the diverse anthropogenic impacts on our planet, with topics ranging from sea-level rise to livestock populations, greenhouse gas emissions, fertilizer use, and beyond. Each entry in the database reports a quantitative value (or a time series of values) along with clear referencing of the primary source, the method of measurement or estimation, an assessment of uncertainty, and links to the underlying data, as well as a permanent identifier called a Human Impacts ID (HuID). While there are other databases that house some of these values, they are typically focused on a single topic area, like energy usage or greenhouse gas emissions. The Human Impacts Database facilitates access to carefully curated data, acting as a quantitative resource pertaining to the myriad ways in which humans have an impact on the Earth, for practicing scientists, the general public, and those involved in education for sustainable development alike. We outline the structure of the database, describe our curation procedures, and use this database to generate a graphical summary of the current state of human impacts on the Earth, illustrating both their numerical values and their intimate interconnections.



INTRODUCTION

One of the most important scientific developments of the last two centuries is the realization that the evolution of Earth is deeply intertwined with the evolution of life. Perhaps the most famous example of this intimate relationship is the large-scale oxygenation of Earth's atmosphere following the emergence of photosynthesis.¹ This dramatic change in the composition of the atmosphere is believed to have caused a massive extinction, as the biosphere was not adapted to an oxygenated atmosphere.^{2–4} Over the past 10,000 years, humans have likewise transformed the planet, directly affecting the rise and fall of ecosystems,^{5–13} the pH and surface temperature of the oceans,^{14,15} the composition of terrestrial biological and human-made mass,^{16,17} the planetary albedo and ice cover,^{18–27} and the chemistry of the atmosphere,^{28–33} to name just a few examples. The breadth of human impacts on the planet is so diverse that it touches on nearly every facet of the Earth system and every scientific discipline.

Technological advances in remote sensing, precision measurement, and computational power have made it possible to measure these anthropogenic impacts with unprecedented depth and resolution. However, as scientists with different training use distinct methods for measurement and analysis, report data in different units and formats, and use nomenclature differently, these studies can be very challenging to understand and relate to one another. Even seemingly simple questions such as “how much water do humans use?” can be difficult to answer when search engines are not optimized for finding numeric data, and a search of the scientific literature yields an array of complicated analyses with different units, varying definitions about what constitutes water use, and distinct approaches to quantifying flows. This problem persists beyond the primary scientific literature, as governmental, intergovernmental, and industry datasets can be similarly tricky and laborious to interpret.

Writing from California, as several of the authors are, where we now have a “wildfire season” and a multi-decadal drought,^{34,35} we wanted to develop a deeper understanding of the ways in which human activities might have produced such dramatic and consequential changes in our local and global environment. In pursuit of basic understanding, we asked many questions, like “how much water and land do humans use?” and “how much methane is emitted annually?” In our search for answers, even when the question is well defined (as is the case for methane emissions), we often encountered the same challenges: disparate technical studies written for expert audiences must be understood, evaluated, and synthesized just to answer simple questions. It seemed to us that a referenced compendium of “things we already know,” akin to the *CRC Handbook of Chemistry and Physics*, would be very useful for us and others.

In building the Human Impacts Database, we took inspiration from our previous experience building and using the BioNumbers Database³⁶ (<https://bionumbers.hms.harvard.edu>), a compendium of quantitative values relating to cell and organismal biology. Over the past decade, the BioNumbers Database has become a widely accessed resource that serves not only as an index of biological numbers, but also as a means of finding relevant primary literature, learning about methods of measurement, and teaching basic concepts in cell biology.³⁷

We believe that a centralized, searchable database for quantitative data encompassing the breadth of human impacts on Earth would be similarly transformative for researchers, students, and the interested public. While reading an entry in the Human Impacts Database is not a replacement for reading the primary literature, the database serves as a resource to expedite the process of finding quantitative data and exploring their interconnection. Importantly, we do not put forward projected scenarios or specific policy proposals for combating anthropogenic effects on Earth. However, we are convinced that such proposals should be evaluated in the light of a comprehensive and quantitative understanding of the Earth-human system.

RESULTS

Finding and compiling numbers from scientific literature, governmental and non-governmental reports, and industrial datasets

We have established the Human Impacts Database (<http://anthroponumbers.org>) as a repository for the rapid discovery of quantities describing the Earth-human system. We here provide a more complete description of the database structure, the values it holds, and the stories it tells us about how humans affect the Earth. As of this writing, the database holds > 300 unique and manually curated entries covering a breadth of data sources, including primary scientific literature, governmental and non-governmental reports, and industrial communications. Before it is added to the database and made public, each entry is vetted extensively by the administrators (see [Note S1](#) for detailed curation procedures). Included in each entry is a summary of the method by which it was determined, an assessment of the corresponding uncertainty, and an explicit statement of any known caveats important for interpretation of the data. While these ≈ 300 entries include those we consider to be essential for a quantitative understanding of human impacts on Earth, it is not an exhaustive list. This database will continue to grow and evolve as more data become publicly released, our understanding of the human-Earth system improves, and members of the scientific community suggest values to be added.

[Figure 1](#) shows the Human Impacts Database Entry for perhaps the most emblematic anthropogenic impact: the standing atmospheric CO₂ concentration. The first two components of an entry are the quantity title and its assigned category and subcategory ([Figures 1A](#) and [1B](#)). Primary categorization falls into one of five classes: “land,” “water,” “energy,” “flora & fauna,” and “atmospheric & biogeochemical cycles.” Of course, these categories are broad, and entries can be associated with several categories. For this reason, each entry is also assigned a narrower “subcategory,” such as “agriculture,” “urbanization,” or “carbon dioxide.” While this categorization is not meant to be exhaustive, and many other schemes could be implemented, we found that this organization allowed us to quickly browse and identify quantities of interest.

Following the title and categorization, we report the measured atmospheric CO₂ concentration. This corresponds to the most recent reported measurement, which is, as of this writing, roughly 416 parts per million (ppm) in 2021 ([Figure 1C](#)). Importantly, we report an approximate value for the CO₂ concentration rather than a precise value to many significant digits. While the most

A Atmospheric CO₂ concentration

B Atmospheric & Biogeochemical Cycles Carbon Dioxide

C Value: ≈ 415 parts per million [see this value in other units](#)

D HuID: 81043

E Relevant Year(s): 1964-2021

F Summary: The present carbon dioxide (CO₂) concentration in the atmosphere as measured at the Mauna Loa Observatory. Temporal variations in the CO₂ concentration can be seen as periodic oscillations corresponding to seasonal changes. The carbon dioxide data on Mauna Loa constitute the longest record of direct measurements of CO₂ in the atmosphere. They were started by C. David Keeling of the Scripps Institution of Oceanography in March of 1958 at a facility of the National Oceanic and Atmospheric Administration.

G Method: Uncertainty in measurements is not reported in original source, and it is assumed lower than the monthly variability. The Mauna Loa data are being obtained at an altitude of 3400 m in the northern subtropics, and may not be the same as the globally averaged CO₂ concentration at the surface. The mass of CO₂ is obtained from the concentration using the molar mass of CO₂, 44 g mol⁻¹; the molar mass of air, 29 g mol⁻¹; and the mass of the atmosphere, 5.15 × 10¹⁸ kg.

H Source: [Scripps CO2 Program Primary Mauna Loa CO2 Record. \(2021\)](#)

I Dataset: [Monthly atmospheric CO2 measurements from Mauna Loa Observatory \(monthly_co2_data_processed.csv\)](#)
regional resolution

J Trend:

K Original Data: [Mauna Loa Free Use Policy](#)

L Added By: ilopezgo

Scripps CO₂ Program
Atmospheric CO₂ Data
Primary Mauna Loa CO₂ Record

Station Name	Station Code	Latitude	Longitude	Elevation (m)
Mauna Loa Observatory, Hawaii	MLO	19.5 °N	155.6 °W	3397

```

year,month,date (decimal),Reported value,Concentration
1958,3,1958.203,monthly mean,315.7
1958,4,1958.288,monthly mean,317.45
1958,5,1958.37,monthly mean,317.51
1958,6,1958.455,monthly mean,
1958,7,1958.537,monthly mean,315.86
1958,8,1958.622,monthly mean,314.93
1958,9,1958.707,monthly mean,313.21
1958,10,1958.789,monthly mean,
1958,11,1958.874,monthly mean,313.33
1958,12,1958.956,monthly mean,314.67
1959,1,1959.041,monthly mean,315.58
1959,2,1959.126,monthly mean,316.49
1959,3,1959.203,monthly mean,316.65
1959,4,1959.288,monthly mean,317.72
    
```

Data sharing policy
The data and graphics on this website are made freely available, with the understanding that appropriate credit will be given. For applications supporting peer-reviewed scientific publications, coauthorship may sometimes be appropriate. An example would be if an important result or conclusion depends on this product, such as the first account of a previously unreported phenomenon. Ethical usage requires disclosing intentions at early stages of the work in order to avoid duplicating ongoing studies at Scripps. For applications where coauthorship is not needed, which includes all applications outside of the peer-reviewed scientific literature, it is sufficient to acknowledge the Scripps CO₂ program as the source. Please direct queries to Ralph Keeling (rkeeling@ucsd.edu)

Figure 1. A representative entry in the Human Impacts Database

(A–I) The entry page for HuID 81043, “Atmospheric CO₂ concentration,” is diagrammed with important features highlighted. Each entry in the Human Impacts Database has (A) a name, (B) a primary and secondary categorization, (C) the numerical value with other units when appropriate, (D) a five-digit permanent numeric identifier, (E) the years for which the measurement was determined, (F) a brief summary of the quantity, (G) the method of determination, (H) a link to the source data, and (I) a link to a processed version of the data saved as a .csv file. When possible, a time series of the data is presented.

(K) Every entry in the database also has a statement of the data use protection associated with the relevant data. When possible, this links directly to the data protection statement from the original source. In other cases, it points to the formal definition of the license by a disinterested third party.

(L) Finally, each entry lists the username of the administrator who curated the quantity. Their contact information is available on the anthronumbers.org “About” page.

recent entry in the linked dataset (Figure 1I) gives a monthly average value of 416.43 ppm for December of 2021, this value does not account for error in the measurement, fluctuations throughout December, or seasonal oscillations in atmospheric CO₂. Therefore, we report a rounded value of 416 ppm. CO₂ measurements are quite accurate, but other measurements and inferences recorded in the Human Impacts Database are less so. We therefore strive to give an assessment of the uncertainty for all values. This can be in the form of a confidence interval, as for the entry for the global mean sea-level rise since 1900 due to thermal expansion, which reports a 90% confidence interval, or bounds on the value, as for the number of contemporary animal extinctions since 1500 CE, which reports only a lower bound. In addition to error assessment, we also aim to provide legible units for all entries. Although atmospheric CO₂ is commonly reported in ppm units, we also report this value in other equivalent units, including the mole and mass fractions of CO₂ and the total mass of CO₂ in the atmosphere in kg CO₂ (Figure 1C). Whenever possible, entries will report values in multiple units to make quantities accessible to readers coming from diverse backgrounds. Furthermore, in many cases, the global value is aggregated from local measurements. We flag entries for which regional data broadly defined are available in the database GitHub repository.

Following the numerical value is the permanent Human Impacts Database identifier, which we abbreviate as HulD (Figure 1D). The HulD is a randomly generated five-digit integer that serves as a permanent and static identifier that can be used for in-line referencing. Rather than identifying a single value, we consider the HulD a pointer to a particular *entry*, so that HulD 81043 can be used to reference the atmospheric CO₂ concentration in 2021 and 1980 (Figure 1E). For example, to reference the present-day atmospheric CO₂ concentration, one could report the value as “≈ 416 ppm (HulD 81043:2021).” In addition, since each entry comes from a single source, we may have more than one HulD reporting similar quantities. For example, HulDs 69674 and 72086 report recent measurements of the temperature of the upper ocean.

The “Summary” field (Figure 1F) gives a succinct description of the quantity and its relationship to “human impacts” broadly construed, along with other pertinent information. This could include a more detailed definition of terms used in the quantity, such as the entry for “sea ice extent loss in March,” which defines the term “sea ice extent,” or useful historical information about the measurement. In our example of atmospheric CO₂ concentration, the summary explains that the measurement is made at the Mauna Loa observatory and points out the seasonal oscillations that are observed. The following “Method” field describes the method by which the quantity was measured, inferred, or estimated (Figure 1G). This field also provides an assessment of the uncertainty in the value, which may include a description of how confidence intervals were computed or a list of critical assumptions that were made to estimate missing data.

All fields through “Method” (Figures 1A–1G) depend on manual curation and interpretation by database administrators. The following two fields, “Source” and “Dataset” (Figures 1H and 1I), provide direct links to the primary source reference and the relevant data. Both of these fields are direct links (shown as insets in Figure 1). The “Source” field can point to either the published scientific literature or the resource page of a govern-

mental, industrial, or non-governmental organization data deposition URL. The “Dataset” field links directly to either a CSV format of the data or to a folder with global and regional values within the corresponding GitHub repository. As discussed in Note S1, the vast majority of these data files have been converted into a “tidy-data” format³⁸ by database administrators, which maximizes programmatic readability.

When possible, a graphical time series of the data is also presented as an interactive plot (Figure 1J). These plots enable users to quickly apprehend time-dependent trends in the data without downloading or processing the dataset. The data sources we rely on in building the database are remarkably varied, coming from governmental, industrial, and primary scientific sources, each with their own specific data use protection policies. Each entry (Figure 1K) also provides a link to the data use policy for each individual dataset. While not available for every entry, the majority of quantities we have curated in the Human Impacts Database contain measurements over time. The last field gives the username of the administrator who generated this entry (Figure 1L). Their affiliation and contact information are available on the database’s “About” page. We invite the reader to contact the administrators collectively—through our “Contact” page or directly through our personal emails as provided on the “About” page—with questions, concerns, or suggestions.

While Figure 1 is a representative example, each quantity in the Human Impacts Database tells a different story. Easy and centralized access to different entries allows users to learn about the magnitude of human impacts and also study the interactions between different human activities, which, as we discuss in the next section, are deeply intertwined.

Global magnitudes

In Figure 2, we provide an array of quantities that we believe to be key in developing a “feeling for the numbers” associated with human impacts on the Earth system. All of the quantities in Figure 2 are drawn from entries in the database and grouped into the same categories used in the database: land, water, flora and fauna, atmosphere and biogeochemical cycles, and energy (see color scheme at the top of Figure 2). Although the impacts considered here necessarily constitute an incomplete description of human interaction with the planet, these numbers encompass many that are critically important, such as the volume of liquid water resulting from ice melt (Figure 2B), the extent of urban and agricultural land use (Figure 2H), global power consumption (Figure 2N), and the heat uptake and subsequent warming of the ocean surface (Figure 2S). In many cases, the raw numbers are astoundingly large and can therefore be difficult to fathom. Rather than reporting only bare “scientific” units, we present each quantity (when possible) in units that are intended to be relatable as “per capita” values to a broad audience who are members of (or familiar with) typical Western lifestyles. Consider, for example, the 18 TW global power consumption (Figure 2N). For most audiences, it can be difficult to conceptualize what a watt is, let alone the sheer magnitude of a *terawatt*. However, most prospective users of this database likely have a familiarity with the warmth of a 100 W light bulb. With this in mind, we can do a simple conversion to say that the global average power use per person is comparable to constantly running ≈ 23 light bulbs per person, making the impact a bit more tangible.




Figure 2. Human impacts on the planet and their relevant magnitudes

Relative units and the broad organizational categories are shown in the top left. Source information and contextual comments for each subpanel are presented in Note S2.

A THE GEOGRAPHY OF HUMAN IMPACTS


Page 1 represents the impact humans have on the Earth at a global scale. While these numbers are handy, it is important to acknowledge that they vary from country-to-country and continent-to-continent. Furthermore, the consequences of these anthropogenic impacts are also unequally distributed, meaning some regions experience effects disproportionate to their contribution. Here, we give a sense of the geographic distribution of several values presented on page 1, broken down by continental region as shown below.



Asia — (As)
North America — (NA)
South America — (SA)
Europe — (Eu)
Oceania — (Oc)
Africa — (Af)


D THE LIVESTOCK POPULATION

The global population of terrestrial livestock is around 30 billion individuals, most of which are chickens. Asia houses most of the global livestock population, though South America and Europe harbor more animals on a per-capita basis.



E NITROGEN FERTILIZER USE & PRODUCTION


Modern agriculture requires nitrogen in amounts beyond what is produced naturally. Asia synthesizes and consumes a large majority of fixed nitrogen. However, Europe and North America dominate per capita synthesis whereas Oceania consumes more fertilizer per capita than any other region.



Source: Food and Agricultural Organization (FAO) of the United Nations. Notes: Values account for reactive nitrogen production in context of fertilizer only and does not account for plastics, explosives, or other uses.

B THE HUMAN POPULATION


There are ≈ 8 billion humans on the planet, with approximately 50% living in 'urban' environments. The majority of the world's population (as well as the majority of both urban and rural dwellers) live in Asia.



Sources: Food and Agricultural Organization of the United Nations - World Population. Notes: Urban/rural designation has no set definition and follows the conventions set by each reporting country.

F WATER WITHDRAWAL


While Asia withdraws the most water for agricultural and municipal needs, North America withdraws the plurality of water for industrial purposes. North America also withdraws more water per capita than any other region.



Source: AQUASTAT Main Database, Food and Agriculture Organization of the United Nations. Notes: Values are reported directly from member countries and represent average of 2013-2017 period. Per capita values are computed given population of reporting countries.

C LAND USE


Though humans are nearly evenly split between urban and rural environments, agricultural land is the far more common use of land area. Together, Asia and Africa contain more than half of global agricultural land. Asia alone accommodates more than half of the global urban land area.



Sources: Food and Agricultural Organization (FAO) of the United Nations (2015) - Land Use (agricultural area). Flörczyk et al. 2019 - GHs Urban Centre Database 2015 (urban land area). Notes: Urban is defined as any inhabited area with ≥ 2500 residents, as defined by the USDA.

G TREE COVERAGE AREA LOSS


Most drivers of tree coverage area loss are comparable in their effect at a global scale. However, there are drastic regional differences in the relative magnitudes.



Source: Curtis et al. 2018 doi: 10.1126/science.aau3445. Notes: Regions are as reported in Curtis et al. 2018. "Deforestation" here denotes permanent removal of tree cover for commodity production. "Shifting agriculture" here denotes forest/shrub land converted to agriculture and later abandoned. All values correspond to breakdown of cumulative tree cover area loss from 2001-2015.

H GREENHOUSE GAS EMISSIONS


CO₂ and CH₄ are two potent greenhouse gases which are routinely emitted by anthropogenic processes such as burning fuel and rearing livestock. While Asia emits roughly half of all CO₂ and CH₄, North America and Oceania produce the most on a per capita basis, respectively.



Sources: CO₂ data collated by: Friedlingstein, P. et al. (2019). doi: 10.5194/essd-11-1783-2019. See Panel K on Pg. 4 for complete list of sources. CH₄ data from Saunori et al. 2020 doi: 10.5194/essd-12-1561-2020. Values report decadal averages in kg CO₂ or CH₄ per year over time period 2008-2017.

I MATERIAL PRODUCTION

Humans excavate an enormous amount of material from the Earth's crust and transform it to build our structures. Two of these materials, concrete and steel, are produced primarily in Asia on both a global and per capita basis. Asia's per capita production of steel is only outpaced by Europe.




Sources: USCS Statistics and Information 2020, Steel Statistical Yearbook 2019 World Steel Association, Food and Agricultural Organization (FAO) of the United Nations - Annual Population. Notes: Reported values for cement and steel production corresponds to 2017 and 2018 values, respectively. Mass of concrete was calculated using a rule-of-thumb that 1 kg of cement yields 7 kg of concrete (Monteiro et al. 2017. doi: 0.138/nmat4930).

J POWER GENERATION AND CONSUMPTION

From heating water, to powering lights, to moving our vehicles, nearly every facet of modern human life requires the consumption of power, culminating in nearly 20 TW of power use in recent years. Asia consumes over half of the power derived from combustion of fossil fuels, with Europe and North America each consuming around 20% of the global total. Asia also produces the plurality of power from renewable technologies, such as hydroelectric, wind, and solar, however, North America, South America, and Europe each produce more on a per capita basis. Nuclear energy, however, is primarily produced in Europe, with North America and Asia coming in second and third place, respectively. On a per-capita basis, North America consumes or produces more energy than all other regions considered here, yielding a total power consumption of nearly 10,000 W per person.

Source: Energy Information Administration of the United States (2017). Notes: "Renewables" includes hydroelectric, biofuels, biomass (wood), geothermal, wind, and solar. "Fossil fuels" includes coal, oil, and natural gas.



(legend on next page)

Exploring these numbers reveals a number of intriguing quantities and relationships. For example, agriculture repeatedly appears as a major contributor to many human impacts, dominating both global land (Figure 2H) and global water use (Figure 2L) and accounting for approximately a third of global tree cover area loss (Figure 2O). In addition, an enormous mass of nitrogen is synthetically fixed through the Haber-Bosch process, primarily to produce fertilizer (Figure 2F), which is a major cause of emissions of N_2O (Figure 2K), which is a potent greenhouse gas. About 45 billion livestock are raised on agricultural lands (Figure 2E), which, together with rice paddies, produce a majority of anthropogenic methane emissions (the greenhouse gas CH_4 ; Figure 2K). On the other hand, urban land area accounts for a very small fraction of land area use ($\approx 1\%$, Figure 2H), and the expansion of cities and suburbs accounts for only $\approx 1\%$ of global tree cover area loss (Figure 2O). This is not to say, however, that urban centers are negligible in their global impacts. As urban and suburban areas currently house more than half of the global human population (Figure 2J), many human impacts are linked to industries that directly or indirectly support urban populations' demand for food, housing, travel, electronics, and other goods. For example, the pursuit of urbanization is the dominating factor in the mass of earth moved on an annual basis (Figure 2W).

Collectively, the ≈ 8 billion humans on Earth (Figure 2J) consume nearly 20 TW of power, equivalent to 23 100 W light bulbs per person (Figure 2N). Around 80% of this energy derives from the combustion of fossil fuels (Figure 2P). This results in a tremendous mass of CO_2 being emitted annually (Figure 2K), of which only $\approx 50\%$ remains in the atmosphere (HuID 70632). A sizable portion of the emissions are absorbed by the oceans (HuID 99089), leading to a steady increase in ocean acidity (Figure 2G) and posing risks to marine ecosystems.³⁹ Furthermore, increasing average global temperatures, primarily caused by greenhouse gas emissions, contribute to sea-level rise not only in the form of added water from melting ice (Figure 2B and 2M), but also due to thermal expansion of ocean water (Figure 2M), which accounts for $\approx 30\%$ of observed sea-level rise.⁴⁰ These are just a few ways in which one can traverse the impacts illustrated in Figure 2, revealing the remarkable extent to which these impacts are interconnected. We encourage the reader to explore this figure in a similar manner, blazing their own trail through the values.

Regional distribution

While Figure 2 presents the magnitude of human impacts at a global scale, it is important to recognize that these impacts—both their origins and their repercussions—are variable across the globe. That is, different societies vary in their preferences for food (e.g., Americans consume relatively little fish) and modes of living (e.g., apartments versus houses), have different levels of economic development (e.g., Canada compared with Malaysia), rely on different natural resources to build infrastructure (e.g., wood versus concrete) and generate power (e.g., nuclear versus coal), and promote different extractive or polluting industries

(e.g., lithium mining versus palm oil farming). Some of these regional differences are evident in Figure 3, which summarizes regional breakdowns of several drivers of global human impacts, e.g., livestock populations and greenhouse gas emissions.

Just as impactful human activities like coal power generation and swine farming are more common in some regions than others (Figure 2), the impacts of human activities affect some regions more than others.⁴² Figure 3 displays a coarse regional breakdown of the numbers from Figure 2 for which regional distributions could be determined from the literature. The region definitions used in Figure 3 are similar to the definitions set forth by the Food and Agricultural Organization (FAO) of the United Nations, assigning the semi-continental regions of North America, South America, Africa (including Russia), Asia, and Oceania. Here, we specify both the total contribution of each region and the per-capita value, given the population of that region as of the year(s) in which the quantity was measured.

Much as in the case of Figure 2, interesting details emerge from Figure 3. For example, Asia dominates global agricultural water withdrawal (excluding natural watering via rainfall), using about 62% of the total, while North America takes the lead in industrial water withdrawal. Interestingly, on a per-capita basis, North America withdraws the most water for all uses: agricultural, industrial, and domestic.

North America also emits more CO_2 per capita than any other region, with Oceania and Europe coming second and third, respectively. This disparity can be partially understood by considering the regional distribution of fossil fuel consumption, the dominant source of CO_2 emissions (Figure 3J). While Asia consumes more than half of the total fossil fuel energy, per-capita consumption is markedly lower than in North America, Europe, and Oceania (Figure 3J). Interestingly, the story is different when it comes to methane. Oceania and South America are the largest emitters of anthropogenic CH_4 , mainly due to a standing population of cattle that rivals that of humans in those regions (Figure 3D) and produces this potent greenhouse gas through enteric fermentation.³³ Regional disparities are also apparent in the means of energy production. While consuming only 4% of the total power, South America generates about 14% of the renewable energy. Nuclear power generation, on the other hand, is dominated by North America and Europe, while Oceania, which has a single research-grade nuclear reactor, generates nearly zero nuclear energy.


Investigating the causes of forest loss by geographic region likewise highlights interesting differences. At a global level, all drivers of forest loss are comparable in magnitude, except for urbanization, which accounts for $\approx 1\%$ of total annual tree cover area loss (Figure 2O). Despite comparable magnitudes, different drivers of forest loss have different long-term consequences.³⁰ Forest loss due to wildfires and forestry often result in regrowth, while commodity-driven harvesting and urbanization tend to be drivers of long-lasting deforestation.^{43,44} Central and South America account for about 65% of commodity-driven deforestation

Figure 3. Regional distribution of anthropogenic effects

(A) Several quantities from Figure 2 were selected, and the relative magnitudes were broken down by subcontinental area. (B–J) Donut charts in all sections show the relative contributions of each quantity by region. Ball-and-stick plots show the per-capita breakdown of each quantity across geographic regions. All data for global and per-capita breakdowns correspond to the latest year for which data were available. The regional breakdown for deforestation uses the regional convention as reported in the source data.⁴¹

A THE HUMAN POPULATION


The human population has more than doubled in the past 60 years. During this time, the fraction of the population living in urban areas has steadily increased such that the global population is about evenly split between urban and rural environments.



Sources: Food and Agricultural Organization of the United Nations - World Population
Notes: Urban/rural designation has no set definition and follows the conventions set by each reporting country.

B WATER WITHDRAWAL


Total water withdrawal has increased in concert with the human population, dominated by increasing agricultural use. Despite this increase, the average per-capita water use for agricultural and domestic purposes has remained largely constant for the past 40 years.




Source: AQUASTAT Main Database, Food and Agriculture Organization of the United Nations.
Notes: Values are reported directly from member countries and represent average of 2013-2017 period. Per capita values are computed given population of reporting countries.

C THE LIVESTOCK POPULATION

The standing population of livestock has been increasing, with chicken making up a large fraction of the total livestock population. The number of chicken raised per capita has increased since the 1960s, while cattle per capita have decreased.






LIVESTOCK PER CAPITA



Sources: Food and Agricultural Organization of the United Nations

D MATERIAL PRODUCTION


The total mass of human-made materials has been accumulating over time, dominated by construction materials. Per capita, the mass of bricks & asphalt, aggregates, and concrete has dramatically increased since the 1950s.



Sources: Krausmann et al. 2017 doi: 10.1073/pnas.1613771114
Notes: Material production is estimated from a material flows model.

E AQUATIC FOODS PRODUCTION


Aquatic (blue) foods production has been increasing with the human population. Interestingly, the mass produced from wild capture has remained constant per capita since the 1980s while the mass produced by aquaculture has increased per capita during the same period, driving the increase in overall production.



Sources: Food and Agricultural Organization of the United Nations

F POWER CONSUMPTION


Power consumption has increased with population, as well as technological and societal changes, which have driven an increase in power per capita across all generation types. The source of our power has also changed over time. Over the last 60 years, nuclear power has become comparable to hydroelectricity, with most of the growth occurring between 1970 and 1990. Renewable power generation is currently experiencing a similar growth pattern.



Source: Energy Information Administration of the United States (2017).
Notes: "Renewables" includes biofuels, biomass (wood), geothermal, wind, and solar. "Fossil fuels" includes coal, oil, and natural gas.


G CO₂ EMISSIONS

Annual anthropogenic CO₂ emissions have been increasing with the population, driven by an increase in fossil fuel combustion. The amount of CO₂ emissions from fossil fuels has increased slightly per capita, while the per capita emissions from land use change have decreased.



H CH₄ EMISSIONS

While total anthropogenic methane (CH₄) emissions have been increasing with the human population, per capita emissions have been decreasing each decade since the 1970s. This per capita reduction reflects a shift in global diets away from methane-intensive beef products, as well as better waste management policies in developed countries.



(legend on next page)

(meaning clear-cutting and human-induced fires with no substantial regrowth of tree cover), whereas a majority of forest loss due to shifting agriculture occurs in Africa (where regrowth does occur). Together, wildfires in North America, Russia, China, and South Asia make up nearly 90% of losses due to fire.⁴¹ While urbanization is the smallest driver of tree cover loss globally, it can still have strong impacts at the regional level, perturbing local ecosystems and biodiversity.^{45,46}

Time series

When available, the Human Impacts Database includes time-series data for each quantity. Just as the regional distributions of impactful human activities help us understand differences between societies and regions, studying the history of these activities highlights recent technological and economic developments that intensify or reduce their impacts. When considering the history of human impacts on the Earth, it is natural to start by considering the growth of the human population over time. As shown in Figure 4, the global human population grew nearly continually over the past 80 years, with the current population nearing 8 billion. Historically, most of the global human population lived in rural areas (about 70% as of 1950, HuID 93995). Recent decades have been marked by a substantial shift in how humans live globally, with around half of the human population now living in urban or suburban settings ($\approx 55\%$, HuID 93995).

Given the growth of the human population, it is reasonable to consider that human population may be the most natural scale to measure human impacts.⁴⁷ To assess this possibility, we plotted per-capita impacts over several decades (Figure 4). If impacts are growing in direct proportion to the human population, per-capita impacts would be constant over time. Indeed, this is roughly true for per-capita water withdrawals over the past 40 years (Figure 4B). Deviations from proportionality may indicate important changes in human activities. For example, in recent decades, per-capita chicken populations grew by nearly 2-fold, while per-capita cattle populations shrunk by roughly 25%, reflecting a modest transition away from beef and toward chicken as a source of animal protein in global diets (HuIDs 40696 and 79776).

One very visible impact accompanying the shift of the human population to urban environments is the increase in production of anthropogenic mass: materials such as concrete, steel, lumber, and plastics used to build roads, buildings, machines, packaging, and other useful human-made items. Since these materials are degraded very slowly, anthropogenic mass has been accumulating over time. In addition, the mass of concrete, aggregates like asphalt, and bricks per capita has been increasing since the 1950s (Figure 4D). Concrete, in particular, has increased from less than 10 tons per person in the 1950s to almost 30 tons per person in the 2010s. This increase in per-capita anthropogenic mass means that the increase in production of these materials is outpacing the growth of the human population.

These material production trends have been enabled, in part, by a sustained increase in power generation. As evident from Figure 4, total power consumption has been increasing roughly proportionally with the human population. Per-capita consumption has also increased across all generation types, including fossil fuels, hydropower, nuclear, and renewables. The growth among nuclear and renewables has been especially dramatic, and nuclear power now roughly equals hydropower production. Production of crops, aquaculture, and populations of livestock are all likewise correlated with growth in the human population (Figures 4C and 4E). The total number of livestock has increased with the human population, primarily due to increasing chicken populations as discussed above. The dominant means of aquatic food production has also shifted over this time: until roughly 1980, nearly all seafood was captured wild, but since then aquaculture has grown to account for roughly $\frac{1}{2}$ of aquatic food production (HuID 61233, Figure 4E).

Turning our focus to greenhouse gases, we see that annual anthropogenic CO₂ emissions have been increasing with the population (Figure 4G). Burning of fossil fuels is the dominant contributor to anthropogenic emissions and has increased slightly on a per-capita basis over the past 60 years. In contrast, as the pace of global deforestation has slowed,^{48,49} emissions of CO₂ due to land-use change have decreased per capita. These two trends roughly neutralize each other, leading to little overall change in CO₂ emissions per capita since the 1960s. Akin to CO₂ emissions due to land-use change, CH₄ emissions show a sub-linear trend with human population, partially due to a decline in ruminant livestock per capita (Figures 4C and 4H).

DISCUSSION

Quantitative literacy is necessary for “understanding” in nearly all branches of science. As our collective knowledge of anthropogenic impacts expands, it has become challenging to sift through the literature to collect specific numbers useful for both calculation and communication. We have attempted to reduce this barrier to entry on several fronts. We have canvassed the scientific literature, governmental, industrial, and international reports to assemble a broad, quantitative picture of how human activities have affected the Earth’s atmosphere, oceans, rivers, lands, biota, chemistry, and geology. In doing so, we have created an online, searchable database housing an array of quantities and data that describe different facets of the human-Earth interface. We view this database as an accessory, rather than a replacement, for the myriad scientific databases that exist and are publicly available on the internet (some of which are listed on the database website www.anthroponumbers.org/catalog/databases). While these databases are invaluable resources for accessing scientific data, the Human Impacts Database is built from the ground up with the intention of being broadly accessible to scientists and the curious general public alike to help build the collective quantitative literacy of the Anthropocene. Beyond the database, we have assembled

Figure 4. Temporal dynamics of key human impacts

(A) Several quantities from Figure 2 were selected, and the magnitudes were plotted as a function of either time (for cumulative quantities such as anthropomass) or human population.

(B–H) Ball-and-stick plots show the per-capita breakdown as decadal averages to give a more reflective view of cultural and technological shifts than year-to-year variation.

these data into a comprehensive snapshot, released alongside this writing as a standalone graphical document (Data S1), with all underlying data, associated uncertainties, and referencing housed in the Human Impacts Database. While necessarily incomplete, these resources provide a broad view of the ways in which human activities are having an impact on the Earth on multiple fronts.

One insight that emerges from a holistic consideration of these diverse human activities together is that they are deeply intertwined and driven by a small number of pivotal factors: the size of the human population, the composition of our diets, and our demand for materials and energy to build and power our increasingly complex and mechanized societies. Understanding the scale of human agriculture and water and power usage provides a framework for understanding most of the numerical gallery presented in Figure 2. Perhaps unsurprisingly, we find that feeding the growing human population is a major driver of a large swath of human impacts on Earth, dominating global land (Figure 2H, HuID 29582) and water use (Figure 2L; HuIDs 84545, 43593, 95345), as well as significantly contributing to tree cover loss (Figure 2O, HuID 24388), earth moving (Figure 2W; HuIDs 19415, 41496), and anthropogenic nitrogen fixation (Figure 2F; HuIDs 60580, 61614), to name a few such examples. The Human Impacts Database provides a resource to explore relationships between values temporally, globally, and locally, and go beyond the standalone values often reported in isolation or cast solely through the lenses of impact, population, affluence, and technology (I = PAT) relationships.

It is common in this setting to argue that the bewildering breadth and scale of human impacts should motivate some specific remediation at the global or local scale. We, instead, take a more modest "just the facts" approach. The numbers presented here show that human activities affect our planet to a large degree in many different and incommensurate ways, but they do not provide a roadmap for the future. Rather, we contend that any plans for the future should be made in the light of a comprehensive and quantitative understanding of the interconnected ways in which human activities impact the Earth system globally (Figure 2), locally (Figure 3), and temporally (Figure 4). Achieving such an understanding will require the synthesis of a broad literature across many disciplines. While the quantities we have chosen to explore are certainly not exhaustive, they represent some of the key axes that frequently drive scientific and public discourse and shape policy across the globe.

Earth is the only habitable planet we know of, so it is crucial to understand how we got here and where we are going. That is, how (and why) have human impacts changed over time? How are they expected to change in the future? For every aspect of human entanglement with the Earth system—from water use to land use, greenhouse gas emissions, mining of precious minerals, and so on—there are excellent studies measuring impacts and predicting their future trajectories. Of particular note are the data-rich and explanatory reports from the Intergovernmental Panel on Climate Change^{50,51} and the efforts toward defining "planetary boundaries."⁵² We hope that the Human Impacts Database and the associated resources with this work provide a reference to explore the human-induced interdependencies between many axes of the human–Earth system and will engage the scientific community, ultimately helping humanity coexist stably with the only planet we have.

EXPERIMENTAL PROCEDURES

Resource availability

Lead contact

Requests for further information should be directed to and will be fulfilled by the lead contact, Griffin Chure (griffinchure@gmail.com).

Materials availability

No materials were used in the generation of this work, other than the code and data as described below. We have collated all data shown in Figures 2–4, along with all information in Note S2 as a printable, "graphical snapshot" (Data S1).

Data and code availability

For every dataset included in the database, there is a folder in the GitHub repository https://github.com/rpgroup-pboc/human_impacts (DOI: 10.5281/zenodo.4453276) that includes the source data, the processed data, and the code to generate the "tidy" data from the source data. Each folder also includes a README file that includes information about the dataset. In addition, all of the code used to generate the figures can be found in the GitHub repository under the "figures" folder. We strongly encourage the scientific community to fork this repository, submit pull requests, and open new constructive issues through the GitHub repository interface.

The database and the FAIR principles of data reuse

The primary goal of the Human Impacts Database is to provide a resource for the rapid discovery quantities related to the human–Earth system while minimizing the grunt work needed to access (and understand) the underlying data. This means that facilitating data reuse and reproducibility of any analyses is paramount to the importance of the database. To that end, we abide by the FAIR Guiding Principles for Scientific Data Management and Stewardship (www.go-fair.org/fair-principles/). These principles are guidelines to maximize the findability, accessibility, interoperability, and reusability of original scientific data. The database closely follows these principles, as is briefly outlined below:

- **Findability:** The underlying data can be easily searched and navigated, permitting rapid discovery. Individual entries are assigned a unique integer identifier that serves as a permanent referencing tool and are provided with rich metadata about the method of determination, original source, data use protection policy, and quantitative value in diverse units.
- **Accessibility:** The original source of the underlying data is always reported hyperlinked when legally permissible. The transformation, collation, or manipulation of the underlying data that was necessary to add it to the Human Impacts Database is preserved under a publicly accessible, version-controlled, GitHub repository (github.com/rpgroup-pboc/human_impacts) and is permanently accessible via <https://doi.org/10.5281/zenodo.4453276>. This protects against permanent loss of the data even if an entry is deleted from the database.
- **Interoperability:** The data are provided in a human readable format with an emphasis on description of the data and their source. The vast majority of datasets are transformed programmatically to follow a "tidy," long-form format that facilitates computational analysis of the data. As the values are hand curated and the target audience is a curious human, we have not developed an API for programmatic access of the database, and do not have plans to do so in the foreseeable future.
- **Reusability:** All entries in the database and the corresponding GitHub repository are extensively annotated with rich metadata, preventing the need for guesswork as to how the data were collected or what the column names refer to in the original or processed data. Furthermore, all data held in the database and repository follow the legal guidelines as presented by their original owner. This licensing is directly linked to in each entry.

SUPPLEMENTAL INFORMATION

Supplemental information can be found online at <https://doi.org/10.1016/j.patter.2022.100552>.

ACKNOWLEDGMENTS

We are incredibly grateful for the generosity of a wide array of experts for their advice, suggestions, and criticism of this work. Specifically, we thank Suzy Beeler, Joseph Berry, Lars Bildsten, Justin Bois, Chris Bowler, Matthew Burgess, Ken Caldeira, Jörn Callies, Sean B. Carroll, Ibrahim Cissé, Joel Cohen, Michelle Dan, Bethany Ehlmann, Gidon Eshel, Moi Exposito-Alonso, Paul Falkowski, Daniel Fisher, Thomas Frederikse, Neil Fromer, Eric Galbraith, Lea Goentoro, Evan Groover, John Grotzinger, Soichi Hirokawa, Greg Huber, Christina Hueschen, Bob Jaffe, Elizabeth Kolbert, Thomas Lecuit, Raphael Magarik, Jeff Marlow, Brad Marston, Jitu Mayor, Elliot Meyerowitz, Lisa Miller, Dianne Newman, Luke Oltrogge, Nigel Orme, Victoria Orphan, Marco Pasti, Pietro Perona, Noam Prywes, Stephen Quake, Hamza Raniwala, Manuel Razo-Mejia, Thomas Rosenbaum, Benjamin Rubin, Alex Rubinsteyn, Shyam Saladi, Tapio Schneider, Murali Sharma, Alon Shepon, Arthur Smith, Matthieu Talpe, Wati Taylor, Julie Theriot, Tadashi Tokieda, Cat Triandifillou, Sabah Ul-Hasan, Tine Valencic, Ned Wingreen, and Emily Zakem. We also thank Yue Qin for sharing data related to global water consumption. Many of the topics in this work began during the Applied Physics 150C course taught at Caltech during the early days of the COVID-19 pandemic. This work was supported by the Resnick Sustainability Institute at Caltech and the Schwartz-Reisman Collaborative Science Program at the Weizmann Institute of Science. G.C. acknowledges support by the NSF Postdoctoral Research Fellowships in Biology Program (grant no. 2010807).

AUTHOR CONTRIBUTIONS

Conceptualization, G.C., R.A.B., R.M., and R.P.; investigation, G.C., R.A.B., A.I.F., N.S.S., M.K., I.L.G., and Y.M.B.; data curation, G.C., R.A.B., N.S.S., M.K., and I.L.G.; software, G.C.; writing – original draft, G.C., R.A.B., A.I.F., N.S.S., I.L.G., R.M., and R.P.; writing – review & editing, G.C., R.A.B., and R.P.; visualization, G.C. and R.A.B.; project administration, G.C., R.A.B., and R.P.

DECLARATION OF INTERESTS

The authors declare no competing interests.

Received: March 4, 2022

Revised: May 26, 2022

Accepted: June 23, 2022

Published: August 3, 2022

WEB RESOURCES

BioNumbers Database, <https://bionumbers.hms.harvard.edu>
Human Impacts Database, <http://anthroponumbers.org>

REFERENCES

- Fischer, W.W., Hemp, J., and Johnson, J.E. (2016). Evolution of oxygenic photosynthesis. *Annu. Rev. Earth Planet Sci.* *44*, 647–683. <https://doi.org/10.1146/annurev-earth-060313-054810>.
- Hodgskiss, M.S.W., Crockford, P.W., Peng, Y., Wing, B.A., and Horner, T.J. (2019). A productivity collapse to end Earth's Great Oxidation. *Proc. Natl. Acad. Sci. USA* *116*, 17207–17212. <https://doi.org/10.1073/pnas.1900325116>.
- Gumsley, A.P., Chamberlain, K.R., Bleeker, W., Söderlund, U., de Kock, M.O., Larsson, E.R., and Bekker, A. (2017). Timing and tempo of the great oxidation event. *Proc. Natl. Acad. Sci. USA* *114*, 1811–1816. <https://doi.org/10.1073/pnas.1608824114>.
- Sessions, A.L., Doughty, D.M., Welander, P.V., Summons, R.E., and Newman, D.K. (2009). The continuing puzzle of the great oxidation event. *Curr. Biol.* *19*, R567–R574. <https://doi.org/10.1016/j.cub.2009.05.054>.
- Estes, J.A., Terborgh, J., Brashares, J.S., Power, M.E., Berger, J., Bond, W.J., Carpenter, S.R., Essington, T.E., Holt, R.D., Jackson, J.B.C., et al. (2011). Trophic downgrading of planet earth. *Science* *333*, 301–306. <https://doi.org/10.1126/science.1205106>.
- Springer, A.M., Estes, J.A., van Vliet, G.B., Williams, T.M., Doak, D.F., Danner, E.M., Forney, K.A., and Pfister, B. (2003). Sequential megafaunal collapse in the North Pacific Ocean: an ongoing legacy of industrial whaling? *Proc. Natl. Acad. Sci. USA* *100*, 12223–12228. <https://doi.org/10.1073/pnas.1635156100>.
- Hale, S.L., and Koprowski, J.L. (2018). Ecosystem-level effects of keystone species reintroduction: a literature review: effects of keystone species reintroduction. *Restor. Ecol.* *26*, 439–445. <https://doi.org/10.1111/rec.12684>.
- Holdo, R.M., Sinclair, A.R.E., Dobson, A.P., Metzger, K.L., Bolker, B.M., Ritchie, M.E., et al. (2009). A disease-mediated trophic cascade in the Serengeti and its implications for ecosystem. *PLoS Biol.* *7*, e1000210. <https://doi.org/10.1371/journal.pbio.1000210>.
- Ceballos, G., Ehrlich, P.R., Barnosky, A.D., García, A., Pringle, R.M., and Palmer, T.M. (2015). Accelerated modern human-induced species losses: entering the sixth mass extinction. *Sci. Adv.* *1*, e1400253. <https://doi.org/10.1126/sciadv.1400253>.
- Davidson, A.D., Hamilton, M.J., Boyer, A.G., Brown, J.H., and Ceballos, G. (2009). Multiple ecological pathways to extinction in mammals. *Proc. Natl. Acad. Sci. USA* *106*, 10702–10705. <https://doi.org/10.1073/pnas.0901956106>.
- Fortin, D., Beyer, H.L., Boyce, M.S., Smith, D.W., Duchesne, T., and Mao, J.S. (2005). Wolves influence Elk movements: Behavior shapes a trophic cascade in Yellowstone National Park. *Ecology* *86*, 1320–1330. <https://doi.org/10.1890/04-0953>.
- Tronstad, L.M., Hall, R.O., Koel, T.M., and Gerow, K.G. (2010). Introduced lake trout produced a four-level trophic cascade in Yellowstone Lake. *Trans. Am. Fish. Soc.* *139*, 1536–1550. <https://doi.org/10.1577/t09-151.1>.
- Palumbi, S.R. (2001). Humans as the world's greatest evolutionary force. *Science* *293*, 1786–1790. <https://doi.org/10.1126/science.293.5536.1786>.
- Balmaseda, M.A., Trenberth, K.E., and Källén, E. (2013). Distinctive climate signals in reanalysis of global ocean heat content. *Geophys. Res. Lett.* *40*, 1754–1759. <https://doi.org/10.1002/grl.50382>.
- Cheng, L., Trenberth, K.E., Fasullo, J., Boyer, T., Abraham, J., and Zhu, J. (2017). Improved estimates of ocean heat content from 1960 to 2015. *Sci. Adv.* *3*, e1601545. <https://doi.org/10.1126/sciadv.1601545>.
- Bar-On, Y.M., Phillips, R., and Milo, R. (2018). The biomass distribution on Earth. *Proc. Natl. Acad. Sci. USA* *115*, 6506–6511. <https://doi.org/10.1073/pnas.1711842115>.
- Elhacham, E., Ben-Uri, L., Grozovski, J., Bar-On, Y.M., and Milo, R. (2020). Global human-made mass exceeds all living biomass. *Nature* *588*, 442–444. <https://doi.org/10.1038/s41586-020-3010-5>.
- IPCC (2019). *The Ocean and Cryosphere in a Changing Climate*.
- Peng, H., Ke, C., Shen, X., Li, M., and Shao, Z. (2020). Summer albedo variations in the Arctic Sea ice region from 1982 to 2015. *Int. J. Climatol.* *40*, 3008–3020. <https://doi.org/10.1002/joc.6379>.
- Marcianesi, F., Aulicino, G., and Wadhams, P. (2021). Arctic sea ice and snow cover albedo variability and trends during the last three decades. *Polar Sci* *28*, 100617. <https://doi.org/10.1016/j.polar.2020.100617>.
- Mouginot, J., Rignot, E., Björk, A.A., van den Broeke, M., Millan, R., Morlighem, M., Noël, B., Scheuchl, B., and Wood, M. (2019). Forty-six years of Greenland Ice Sheet mass balance from 1972 to 2018. *Proc. Natl. Acad. Sci. USA* *116*, 9239–9244. <https://doi.org/10.1073/pnas.1904242116>.
- MacGregor, J.A., Fahnestock, M.A., Catania, G.A., Aschwanden, A., Clow, G.D., Colgan, W.T., Gogineni, S.P., Morlighem, M., Nowicki, S.M.J., Paden, J.D., et al. (2016). A synthesis of the basal thermal state of the Greenland Ice Sheet. *J. Geophys. Res. Earth Surf.* *121*, 1328–1350. <https://doi.org/10.1002/2015jf003803>.
- Kwok, R. (2018). Arctic sea ice thickness, volume, and multiyear ice coverage: losses and coupled variability (1958–2018). *Environ. Res. Lett.* *13*, 105005. <https://doi.org/10.1088/1748-9326/aae3ec>.
- Massom, R.A., Scambos, T.A., Bennetts, L.G., Reid, P., Squire, V.A., and Stammerjohn, S.E. (2018). Antarctic ice shelf disintegration triggered by

- sea ice loss and ocean swell. *Nature* 558, 383–389. <https://doi.org/10.1038/s41586-018-0212-1>.
25. Notz, D., and Stroeve, J. (2016). Observed Arctic sea-ice loss directly follows anthropogenic CO₂ emission. *Science* 354, 747–750. <https://doi.org/10.1126/science.aag2345>.
 26. Stroeve, J., and Notz, D. (2018). Changing state of Arctic sea ice across all seasons. *Environ. Res. Lett.* 13, 103001. <https://doi.org/10.1088/1748-9326/aade56>.
 27. Goode, P.R., Pallé, E., Shoumko, A., Shoumko, S., Montañes-Rodríguez, P., and Koonin, S.E. (2021). Earth's albedo 1998–2017 as measured from earthshine. *Geophys. Res. Lett.* 48. <https://doi.org/10.1029/2021gl094888>.
 28. Friedlingstein, P., Jones, M.W., O'sullivan, M., Andrew, R.M., Hauck, J., Peters, G.P., Peters, W., Pongratz, J., Sitch, S., Le Quéré, C., et al. (2019). Global carbon budget 2019. *Earth Syst. Sci. Data* 11, 1783–1838.
 29. Houghton, R.A., and Nassikas, A.A. (2017). Global and regional fluxes of carbon from land use and land cover change 1850–2015: Carbon emissions from land use. *Glob. Biogeochem. Cycles* 31, 456–472. <https://doi.org/10.1002/2016gb005546>.
 30. Hansis, E., Davis, S.J., and Pongratz, J. (2015). Relevance of methodological choices for accounting of land use change carbon fluxes. *Glob. Biogeochem. Cycles* 29, 1230–1246. <https://doi.org/10.1002/2014gb004997>.
 31. Tian, H., Xu, R., Canadell, J.G., Thompson, R.L., Winiwarter, W., Suntharalingam, P., Davidson, E.A., Ciais, P., Jackson, R.B., Janssens-Maenhout, G., et al. (2020). A comprehensive quantification of global nitrous oxide sources and sinks. *Nature* 586, 248–256. <https://doi.org/10.1038/s41586-020-2780-0>.
 32. Keeling, C.D. (1960). The concentration and isotopic abundances of carbon dioxide in the atmosphere. *Tellus* 12, 200–203. <https://doi.org/10.1111/j.2153-3490.1960.tb01300.x>.
 33. Saunois, M., Stavert, A.R., Poulter, B., Bousquet, P., Canadell, J.G., Jackson, R.B., Raymond, P.A., Dlugokencky, E.J., Houweling, S., Patra, P.K., et al. (2020). The global methane budget 2000–2017. *Earth Syst. Sci. Data* 12, 1561–1623.
 34. Yoon, J.-H., Kravitz, B., Rasch, P.J., Simon Wang, S.Y., Gillies, R.R., and Hippias, L. (2015). Extreme fire season in California: A glimpse into the future? *Bull. Am. Meteorol. Soc.* 96, S5–S9. <https://doi.org/10.1175/bams-d-15-00114.1>.
 35. Seager, R., Hoerling, M., Schubert, S., Wang, H., Lyon, B., Kumar, A., Nakamura, J., and Henderson, N. (2015). Causes of the 2011–14 California drought. *J. Clim.* 28, 6997–7024. <https://doi.org/10.1175/jcli-d-14-00860.1>.
 36. Milo, R., Jorgensen, P., Moran, U., Weber, G., and Springer, M. (2010). BioNumbers—the database of key numbers in molecular and cell biology. *Nucleic Acids Res.* 38, D750–D753. <https://doi.org/10.1093/nar/gkp889>.
 37. Milo, R., and Phillips, R. (2016). *Cell Biology by the Numbers* (Garland Science).
 38. Wickham, H. (2014). Tidy data. *J. Stat. Softw.* 59. <https://doi.org/10.18637/jss.v059.i10>.
 39. Andersson, A., Kline, D., Edmunds, P., Archer, S., Bednaršek, N., Carpenter, R., Chadsey, M., Goldstein, P., Grottolli, A., Hurst, T., et al. (2015). Understanding ocean acidification impacts on organismal to ecological scales. *Oceanography* 25, 16–27. <https://doi.org/10.5670/oceanog.2015.27>.
 40. Frederikse, T., Landerer, F., Caron, L., Adhikari, S., Parkes, D., Humphrey, V.W., Dangendorf, S., Hogarth, P., Zanna, L., Cheng, L., and Wu, Y.H. (2020). The causes of sea-level rise since 1900. *Nature* 584, 393–397. <https://doi.org/10.1038/s41586-020-2591-3>.
 41. Curtis, P.G., Slay, C.M., Harris, N.L., Tyukavina, A., and Hansen, M.C. (2018). Classifying drivers of global forest loss. *Science* 361, 1108–1111. <https://doi.org/10.1126/science.aau3445>.
 42. Patz, J.A., Campbell-Lendrum, D., Holloway, T., and Foley, J.A. (2005). Impact of regional climate change on human health. *Nature* 438, 310–317. <https://doi.org/10.1038/nature04188>.
 43. Bowman, D.M.J.S., Balch, J.K., Artaxo, P., Bond, W.J., Carlson, J.M., Cochrane, M.A., D'Antonio, C.M., DeFries, R.S., Doyle, J.C., Harrison, S.P., et al. (2009). Fire in the earth system. *Science* 324, 481–484. <https://doi.org/10.1126/science.1163886>.
 44. Santín, C., Doerr, S.H., Preston, C.M., and González-Rodríguez, G. (2015). Pyrogenic organic matter production from wildfires: A missing sink in the global carbon cycle. *Glob. Change Biol.* 21, 1621–1633. <https://doi.org/10.1111/gcb.12800>.
 45. Olivier, T., Thébaud, E., Elias, M., Fontaine, B., and Fontaine, C. (2020). Urbanization and agricultural intensification destabilize animal communities differently than diversity loss. *Nat. Commun.* 11, 2686. <https://doi.org/10.1038/s41467-020-16240-6>.
 46. Newbold, T., Hudson, L.N., Hill, S.L.L., Contu, S., Lysenko, I., Senior, R.A., Börger, L., Bennett, D.J., Choimes, A., Collen, B., et al. (2015). Global effects of land use on local terrestrial biodiversity. *Nature* 520, 45–50. <https://doi.org/10.1038/nature14324>.
 47. Syvitski, J., Waters, C.N., Day, J., Milliman, J.D., Summerhayes, C., Steffen, W., Zalasiewicz, J., Cearreta, A., Galuszka, A., Hajdas, I., et al. (2020). Extraordinary human energy consumption and resultant geological impacts beginning around 1950 CE initiated the proposed Anthropocene Epoch. *Commun. Earth Environ.* 1, 32. <https://doi.org/10.1038/s43247-020-00029-y>.
 48. FAO (2020). *State of the World's Forests 2020*.
 49. Minx, J.C., Lamb, W.F., Andrew, R.M., Canadell, J.G., Crippa, M., Döbbeling, N., Forster, P.M., Guizzardi, D., Olivier, J., Peters, G.P., et al. (2021). A comprehensive and synthetic dataset for global, regional, and national greenhouse gas emissions by sector 1970–2018 with an extension to 2019. *Earth Syst. Sci. Data* 13, 5213–5252. <https://doi.org/10.5194/essd-13-5213-2021>.
 50. Masson-Delmotte, V., Zhai, P., Pörtner, H.O., Roberts, D., Skea, J., Shukla, P.R., et al. (2018). In *Special Report: Global Warming of 1.5 oC* (IPCC).
 51. Pörtner, H.-O., Roberts, D.C., Masson-Delmotte, V., and Zhai, P., eds. (2019). *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate* (IPCC).
 52. Steffen, W., Richardson, K., Rockström, J., Cornell, S.E., Fetzer, I., Bennett, E.M., Biggs, R., Carpenter, S.R., de Vries, W., de Wit, C.A., et al. (2015). Sustainability. Planetary boundaries: guiding human development on a changing planet. *Science* 347, 1259855. <https://doi.org/10.1126/science.1259855>.